

Open clusters in a dispersing molecular cloud

Hui-Chen Chen¹, Chung-Ming Ko^{1,2}

¹ Institute of Astronomy, National Central University, Taiwan

² Department of Physics and Center for Complex System, National Central University, Taiwan

E-mail(HC):huichen@astro.ncu.edu.tw

ABSTRACT

Star clusters are formed in molecular clouds which are believed to be the birth places of most stars. From recent observational data, Lada & Lada(2003) estimated that only 4 to 7% of the proto-clusters have survived. Many factors could cause this high infant mortality. Galactic tidal forces, close encounters with molecular clouds and shock heating are among the possible causes but they have a longer timescale than typical lifetime of molecular clouds. Another possible reason is mass loss in very beginning of cluster evolution in the form of UV radiation, stellar winds or supernova explosions. Mass loss is the main factor we study in this work by using N-body simulations. We find that most proto-clusters survive for more than 40 Myr even when the mass loss rate is high.

1. Introduction

Stellar clusters are among the most interesting objects in astronomy. We are interested in how they form, how they evolve and how they die. Theories and observations indicate that most stars are not born independently but in stellar clusters or stellar associations which are formed in molecular clouds. Due to the limits of observational techniques, we do not know in detail the relation between the clusters and the parent molecular clouds. It is believed that stars are born in clusters and become field stars after the clusters disassociate.

Recently, near infrared observational data (2MASS, Two Micron All Sky Survey project at IPAC/Caltech) have shown that the number of embedded clusters is much higher than the number of optical clusters for which the parent clouds have already dissipated and that the survival probability for proto-clusters is about 4 to 7% (Lada and Lada 2003). This implies that clusters are likely to be disrupted before the clouds are dissipated completely. From the time the clusters are born, the surrounding environments keep dissolving them. Galactic tidal forces, close encounters with giant molecular clouds, shock heating and mass loss by massive member stars (Boily and Kroupa 2003) are possible dissolving mechanisms that operate during the clusters' lifetime. Nonetheless, most of them have a longer timescale than the lifetime of molecular clouds, which is less than 10 Myr (Williams et al 1999). Naively speaking, the mechanism which works within the lifetime of the clouds should be the main reason of this low survival probability. In this work, we focus on the effects of mass loss of the cloud in the early evolution of the system.

In the beginning, the clusters are bound to the molecular clouds. As the clouds dissipate, the binding energy

from the cloud decreases and the stellar systems become out of equilibrium. Once out of equilibrium, they may expand or dissociate completely.

2. Simulations

Due to the very different size and density between the clouds and the clusters, it is not easy to simulate stars and gas clouds together. In this work, we adopted the NBODY2 code developed by Aarseth (Aarseth 2001) to study the dynamical evolution of the clusters.

2.1. Model for cloud dispersion

The initial distribution of the clouds is represented by a potential energy in the form of the Plummer model(1911),

$$\Phi_P = \frac{-GM_b}{\sqrt{r^2 + a^2}}, \quad (1)$$

where M_b is the mass of the cloud, a is the length scale of the potential and G is the gravitational constant. To model the dispersion of the cloud we allow the potential energy to evolve in time according to

$$a = a_0^{\alpha t}, \quad (2)$$

where a_0 is the initial length scale of the cloud and α is the dispersion rate of the cloud. The mass of the cloud, M_b remains constant as the length scale increases with time t . The potential well of this system tends to become shallower in time so that the velocity of some stars can exceed the escape velocity and the stars can run away.

2.2. Initial conditions

The stars are initially distributed according to a Plummer distribution both in physical positions and velocities which are required to achieve virial equilibrium. Note that the fact the clusters are in virial equilibrium does not mean that they are also in dynamical equilibrium (Goodwin 1977). The steps for generating the initial conditions are as followed:

- generate a cluster with a Plummer distribution and with a size of about 1 pc.
- put the cluster into a molecular cloud, represented as a Plummer potential energy, in the same center of mass and set the dispersion rate α of the cloud to zero, which means that the molecular cloud will not change with time.
- run the code until the cluster is in a quasi-steady state.
- use this quasi-steady state as initial condition for the simulations, set the dispersion rate α to be greater than zero and let the evolution start.

In our simulations we study three free parameters. First, the mass of the cloud, M_b , is a multiple of the stellar cluster mass M_c , from 1 to 10, resulting in a star formation efficiency,

$$\eta = \frac{M_c}{M_c + M_b}, \quad (3)$$

of 50 to 9%, respectively. Second, the compactness of the cloud, which is controlled by the initial length scale of the potential energy, a_0 , runs from 0.25 pc to 2.5 pc. Third, the dispersion rates of the cloud, α , are 0.05, 0.1 and 0.2 Myr^{-1} . We performed three hundred runs which are described in Table 1 for the single mass cases. Table 2 lists the parameters of three typical cases which we select for further analysis.

In each simulation, we use 2500 particles and consider two kinds of mass function: a single mass distribution and a mass function with slope is -3 (where Salpeter is -2.35) from 0.15 to 12 $M_\odot$.

3. Results

3.1. Case study

In order to investigate parameter space (M_b , a_0), we perform a few hundred runs for this work. To illustrate the main results, we present three typical cases which are listed in Table 2: (1Aa) $M_b=1$ stellar cluster mass, $a_0=2.5$ pc, less massive and loose cloud. (5Ea) $M_b=5$ stellar cluster mass, $a_0=1.25$ pc, intermediate property cloud. (10Ja) $M_b=10$ stellar cluster mass, $a_0=2.5$ pc,

Table 1. parameters

Run	1	2	3	...	8	9	10
$M_b [M_c]$	1	2	3	...	8	9	10
$\eta [\%]$	50	33	25	...	11	10	9
Run	A	B	C	...	H	I	J
$a_0 [\text{pc}]$	0.25	0.5	0.75	...	2	2.25	2.5
Run	a	b	c				
$\alpha [\text{Myr}^{-1}]$	0.2	0.1	0.05				

Table 2. case study

Run	M_b [$M_{cluster}$]	a_0 [pc]	α [Myr^{-1}]
1Aa	1	2.5	0.2
5Ea	5	1.25	0.2
10Ja	10	0.25	0.2

massive and compact cloud. Fig.1 shows the number of stars varying in time while Fig.2 shows the Lagrangian radii (the radii which contain different percentages of the cluster mass) for the three cases. We find that:

Case 1Aa: the masses of the cloud and of the cluster are comparable, which means that the binding energy of this system is not high in the beginning. The structure of the cluster was rarely affected by the dispersion of the cloud which can be seen in Fig.1(a) and fig.2(a). In Fig.1(a), the number of stars decreases slowly in time, and it does not change significantly even after 250 Myr. In the inner region, 0.5 pc, the number of stars increases slightly due to the cluster contraction. Fig.2(a) indicates that only the 90% Lagrangian radius increases by a factor of two while the 50 and 70% Lagrangian radii increase slightly. Similarly, the 10 and 30% Lagrangian radii show a small increase. In this case, the cluster expands in the outer part but contracts in the inner part. More than 2000 stars remain in the inner 2 pc even after 250 Myr. We therefore argue that the cluster is still bound.

Case 5Ea: the star formation efficiency is 13% and the binding energy is much higher than Case 1Aa. The number of stars decrease quickly in the first 5 Myr, but becomes stable after 10 Myr, as shown in Fig.1(b). Fig.2(b) indicates that all the Lagrangian radii increase in the beginning but stop at about 10 Myr, with the except 90% Lagrangian radius, which expands for a longer time but also becomes stable after 35 Myr. Therefore, the cluster

expands due to the dispersion of the cloud and becomes looser: the number density reduces to half within a 2 pc radius.

Case 10Ja: in this very low star formation efficiency case, shown in Fig.1(c) and Fig.2(c), the number of stars drops to zero in 10 Myr and the Lagrangian radii increase. The expansion continues till the end of the simulation. The cluster is disrupted within a few Myr.

3.2. Systems with equal mass

Fig.3(a) shows the final number of stars within 2 pc in the runs with a single mass distribution. In larger and less massive clouds, the final number of stars within 2 pc is high. The number of stars decreases as a_0 becomes smaller and M_b becomes larger. In fig.3(b), the final 50% Lagrangian radius is high for smaller a_0 and larger M_b . The final radius decreases as a_0 becomes larger and M_b becomes smaller. The density for 500 stars in 2 pc radius is about $15/\text{pc}^3$, which is much higher than the field star density, $0.1/\text{pc}^3$. The 500 contour line in the final number plot and the 5 pc contour line in the final radius plot match perfectly. We conclude that the clusters survived at least beyond this line.

3.3. Systems with mass function

Observations show that there is an initial mass function in stellar clusters. Here we generate the clusters by taking -3.0 as the mass function slope (where the Salpeter's slope is -2.35), and the mass range is from 0.15 to $12 M_\odot$. In every simulation, we also consider 2500 stars as in the cases of the single mass model. Fig.4 shows that the trend of the final number of stars within 2 pc and the final radius of the inner 50% of stars are similar to the simulations for the single mass distribution. Also, the 500 contour line in the final number plot and the 5 pc contour line in the final radius plot match each other.

3.4. Dispersion rates

There are three dispersion rates of the cloud we adopted for the single mass model, 0.05, 0.1 and 0.2 Myr^{-1} . It seems that even in the $\alpha=0.05 \text{ Myr}^{-1}$ cases, for which the e-fold time is 4 times smaller than 0.2 Myr^{-1} , the result is quite similar. The larger dispersion rates will speed the evolution, but not much. This supports the idea that the survival probability is high during the cloud dissipation even with a very high dissipation rate.

4. Conclusions

The proto-clusters can be disrupted by mass loss in the early stages of evolution. In this work, we studied this behavior by means of N-body simulations. The mass loss tends to expand or even disassociate the clusters due to the decrease in binding energy. The survival probabilities

are high both when the star formation efficiency is high and when the molecular cloud is loose. More than 75% of the clusters retain a core, with a number density higher than $15/\text{pc}^3$ after few hundred Myr. Different dispersing rates for the cloud provide similar results and even the largest rate does not disrupt all the clusters. Systems with and without the initial mass function have different final densities but agree with each other very well. We conclude that the infant mortality should be low if the proto-clusters are bound from the beginning.

Acknowledgements

The authors appreciate Alessia Gualandris for her kindly and very helpful discussions. This work was supported in part by the National Science Council, Taiwan under the Grants NSC-093-2112-M-008-017 and NSC-94-2112-M-008-018.

References

- Aarseth, S.J. 2001 *New Astronomy*, 6, 277
- Boily, C.M. and Kroupa, P. 2003 *MNRAS*, 338, 665
- Boily, C.M. and Kroupa, P. 2003 *MNRAS*, 338, 673
- Goodwin, S.P. 1997 *MNRAS*, 284, 785
- Lada, C..J. and Lada, E.A. 2003 *ARA&A*, 41, 57




Fig. 1. Number of stars in fix radii at $\alpha=0.2 \text{ Myr}^{-1}$. (a) Case 1Aa, (b) Case 5Ea and (c) Case 10Ja.




Fig. 2. Lagrangian radii at $\alpha=0.2 \text{ Myr}^{-1}$. (a) Case 1Aa, (b) Case 5Ea and (c) Case 10Ja.




Fig. 3. Result for single mass distribution at $\alpha=0.2 \text{ Myr}^{-1}$. (a) final number of stars within 2 pc, (b) final 50 % Lagrangian radius [pc].




Fig. 4. Result for stars with a IMF slope=3.0 at $\alpha=0.2 \text{ Myr}^{-1}$. (a) final number of stars within 2 pc, (b) final 50 % Lagrangian radius [pc].